arXiv:1601.04300v1 [math.DG] 17 Jan 2016

Reductions of Gauss-Codazzi equations

By Robert Conte [1,2] *and A. Michel Grundland* [3,4]

Submitted 22 August 2015, revised 10 December 2015, accepted 14 January 2016

We prove that conformally parametrized surfaces in Euclidean space $\mathbb{R}^3(c)$ of curvature c admit a symmetry reduction of their Gauss-Codazzi equations whose general solution is expressed with the sixth Painlevé function. Moreover, it is shown that the two known solutions of this type (Bonnet 1867, Bobenko, Eitner and Kitaev 1997) can be recovered by such a reduction.

Keywords: Gauss-Codazzi equations; symmetry reduction; sixth Painlevé equation.

PACS

02.20.Sv,
02.30.Hq Ordinary differential equations
02.30.Ik Integrable systems
02.30.Jr Partial differential equations
02.30.+g
02.40.-k Differential geometry
02.40.Hw Classical differential geometry
20.40.Dr

MSC

37K15 Integration of completely integrable systems by inverse spectral and scattering methods
Primary 53A05
Secondary 58F07

Centre de mathématiques et de leurs applications, École normale supérieure de Cachan, 61, avenue du Président Wilson, F–94235 Cachan Cedex, France. Robert.Conte@cea.fr, Grundlan@crm.umontreal.ca

DOI: 10.1146/((please add article doi))
STUDIES IN APPLIED MATHEMATICS AA:1–25
© 2016 Wiley Periodicals, Inc., A Wiley Company

Contents

1 Introduction **3**

2 Previous solutions involving P6 **5**
2.1 Constraint to be a Bonnet surface 6
2.2 Constraint to be a surface dual to a Bonnet surface 8

3 Lie point symmetries and reductions **10**
3.1 Lie point symmetries of the Gauss-Codazzi system 10
3.2 Reduction 11
3.2.1 Reduction, generic isothermic case, $g^2 \neq 1, c = 0$. 12
3.2.2 Reduction, isothermic case, $g = 1, c = 0$ 14
3.2.3 Reduction, isothermic case, $g = -1, c$ arbitrary . . 14
3.3 On the linear representations of P6 17

4 Conclusion **19**

1. Introduction

Consider a three-dimensional Riemannian manifold $\mathbb{R}^3(\kappa)$ having a constant curvature κ. When κ is respectively negative, zero, positive, this three-dimensional manifold is respectively the hyperbolic space $\mathbb{H}^3(\kappa)$, the Euclidean space $\mathbb{R}^3$, the sphere $\mathbb{S}^3(\kappa)$ of radius $\kappa^{-1/2}$. For convenience we will denote $\kappa = -c^2 \in \mathbb{R}$ and also denote the just mentioned spaces $\mathbb{R}^3(c)$ as, respectively, $\mathbb{H}^3(c), \mathbb{R}^3, \mathbb{S}^3(c)$.

Let $\mathbf{F}$ be an immersion of some complex two-dimensional Riemannian manifold into $\mathbb{R}^3(c)$. In conformal coordinates $z = x + iy$, $\bar{z} = x - iy$, the two fundamental forms of the surface are given by

$$\mathrm{I} = < \mathrm{d}\mathbf{F}, \mathrm{d}\mathbf{F} > = e^u \mathrm{d}z \ \mathrm{d}\bar{z}, \tag{1}$$

and

$$\mathrm{II} = - < \mathrm{d}\mathbf{F}, \mathrm{d}\mathbf{N} > = Q\mathrm{d}z^2 + e^u H \mathrm{d}z \ \mathrm{d}\bar{z} + \overline{Q}\mathrm{d}\bar{z}^2, \tag{2}$$

in which $\mathbf{N}$ is a unit vector normal to the tangent plane, u and H are real valued functions, and Q is a complex valued function.

In terms of the principal curvatures $1/R_1$ and $1/R_2$, the mean curvature H and the total (or Gaussian) curvature K are defined as

$$H = \frac{1}{2}\left(\frac{1}{R_1} + \frac{1}{R_2}\right), \tag{3}$$

$$K = \frac{1}{R_1 R_2} = H^2 - c^2 - 4e^{-2u}|Q|^2 = -2e^{-u}u_{z\bar{z}}. \tag{4}$$

The moving frame defined by

$$\sigma = \begin{cases} {}^{\mathrm{t}}(\mathbf{F}_z, \mathbf{F}_{\bar{z}}, \mathbf{N}) & (c = 0), \\ {}^{\mathrm{t}}(\mathbf{F}, \mathbf{F}_z, \mathbf{F}_{\bar{z}}, \mathbf{N}) & (c \neq 0), \end{cases} \tag{5}$$

evolves linearly as

$$\sigma_z = U\sigma, \ \sigma_{\bar{z}} = V\sigma, \tag{6}$$

in which the two linear operators only depend on $u, Q, \overline{Q}, H, c$.

Instead of representing U and V as third ($c = 0$) or fourth ($c \neq 0$) order matrices (or even fifth order ones [1] when $c = 0$), it proves quite convenient to use the representation by second order matrices [2, 3],

$$U = \begin{pmatrix} (1/4)u_z & -Qe^{-u/2} \\ (1/2)(H+c)e^{u/2} & -(1/4)u_z \end{pmatrix}, \ V = \begin{pmatrix} -(1/4)u_{\bar{z}} & -(1/2)(H-c)e^{u/2} \\ \overline{Q}e^{-u/2} & (1/4)u_{\bar{z}} \end{pmatrix} \tag{7}$$

The zero-curvature (Maurer-Cartan) condition

$$[\partial_z - U, \partial_{\bar{z}} - V] = U_{\bar{z}} - V_z + UV - VU = 0, \tag{8}$$

generates the Gauss-Codazzi system of equations,

$$\left\{ \begin{array}{l} u_{z\bar{z}} + \dfrac{1}{2}(H^2 - c^2)e^u - 2|Q|^2 e^{-u} = 0 \ \ \text{(Gauss)}, \\ Q_{\bar{z}} - \dfrac{1}{2} H_z e^u = 0, \ \overline{Q}_z - \dfrac{1}{2} H_{\bar{z}} e^u = 0 \ \text{(Codazzi)}. \end{array} \right. \tag{9}$$

After the initial work of the Polish school [4, 5, 6], who interpreted, among others, isothermic surfaces in $\mathbb{R}^3$ as "soliton surfaces", the relation between geometry and integrability is now well established, see the lecture notes [7] for a summary. Many integrable 1+1-dimensional partial differential equations (PDEs) are contained in the Gauss-Codazzi system (9) [6, 8]. In particular, two solutions have been found, separated by more than one century [9, 10], in terms of the master Painlevé function P6 (Appendix 4) or its confluence to P5. Other solutions have also been found in terms of P3 [10, Eq. (20)].

The motivation of the present paper is the following. On one hand, the Gauss-Codazzi equations completely describe the geometry. On the other hand, the P6 equation is complete in the classical sense [11], i.e. it is impossible to add any term to P6 under penalty of losing the Painlevé property (singlevaluedness of the general solution near all those singularities which depend on the initial conditions). Comparing the total number of arbitrary constants in P6 (six, i.e. two movable constants plus four fixed constants θ_j, see Eq. (A6)) and the small difference (one) between the sum of the differential order of (9) (four) and the number of its parameters (one, c), it is natural to search for a solution in terms of the full P6, i.e. to remove the two constraints among the four θ_j in the two existing P6 solutions [9, 12, 10, 13].

Since surfaces described by Painlevé equations ultimately arise from some ordinary differential equation (ODE), in the present paper we systematically look for reductions of the PDE system (9) to a system of ODEs.

The paper is organized as follows.

In section 2, we recall for later reference the two solutions, found up to now, of the PDE system (9) in terms of P6.

In section 3, we establish the Lie symmetries of (9) and define essentially one reduction to a system of ODEs, the reduced independent variable being $\xi = G_1(z) - G_2(\bar{z})$, with G_1 and G_2 arbitrary. The reduced system, which depends on one additional parameter g, is further integrated with the isothermic constraint, i.e. Q real in some local coordinate. The generic case $g^2 \neq 1$ provides a new solution, in which $e^u H/Q$ and the logarithmic derivatives of H, Q with respect to ξ are P6 functions.

The two nongeneric cases $g = -1$ and $g = 1$ allow us to recover, therefore in a systematic way, the two previous P6 solutions.

2. Previous solutions involving P6

In the point of view adopted here, the system (9) is a set of three coupled partial differential equations (PDEs) in four unknowns $(u, H, Q, \overline{Q})$, and, in contrast to the geometric approach, these four unknowns will be considered as *independent* and *complex*.

Since the Gauss-Codazzi equations are underdetermined, one can impose an additional constraint on the variables of (9). Before presenting these constraints and listing the resulting ODE systems defined in the complex domain, let us recall some invariance properties and give two definitions (Bonnet surface, isothermic surface), or more precisely two characteristic properties.

This nonlinear system (9) admits two types of invariance. The first one is the conformal invariance

$$\begin{aligned}\forall G_1, G_2 :\ & (z, \bar{z}, e^u, H, Q, \overline{Q}) \\ & \to (G_1(z), G_2(\bar{z}), G_1'(z)G_2'(\bar{z})e^u, H, G_1'(z)^2 Q, G_2'(\bar{z})^2 \overline{Q}).\end{aligned} \tag{10}$$

The second one only exists under the condition $Q - \overline{Q} = c$, this is the involution [2, Eq. (4.4)] [13, p. 77] [3, §3 p. 6]

$$(u, H, Q, \overline{Q}) \to \left(-u, 2Q - c = 2\overline{Q} + c, \frac{H + c}{2}, \frac{H - c}{2}\right). \tag{11}$$

THEOREM 1. *[14, Theorem 2.1] A Bonnet surface in $\mathbb{R}^3(c)$ is characterized by the two conditions*

$$(\log Q)_{z\bar{z}} - (\log Q)_{\bar{z}}(\log \overline{Q})_z = 0,\ (\log \overline{Q})_{z\bar{z}} - (\log \overline{Q})_z(\log Q)_{\bar{z}} = 0. \tag{12}$$

This was proven for $c = 0$ in [15] and for any c in [16].

THEOREM 2. *[14, Theorem 3.2] An isothermic surface in $\mathbb{R}^3(c)$ is characterized by the condition*

$$(\log Q - \log \overline{Q})_{z\bar{z}} = 0. \tag{13}$$

Since the relation (13) is a consequence of the two relations (12), any Bonnet surface is isothermic, but not conversely, see an example section 2.2.

The notation c with a subscript ($c_{\mathrm{z}}, c_{\mathrm{u}}, c_{\mathrm{h}}, c_{\mathrm{q}}$, etc) denotes a complex-valued constant.

2.1. Constraint to be a Bonnet surface

Since any Bonnet surface is isothermic, if it is in addition umbilic-free, there exists an isothermic coordinate z such that $1/Q(z,\bar z)$ is harmonic. Let us therefore impose the constraint that $1/Q$ be harmonic,

$$\frac{1}{Q} = f(z) + \bar f(\bar z). \tag{14}$$

This will lead to a solution in terms of P6, first found when $c = 0$ by Bonnet in 1867 [9, p. 84] (although often attributed to Hazzidakis [12, p. 48]), and later extrapolated to arbitrary c by Bobenko and Eitner [13, §4.5].

If instead of z one considers the conformal coordinate w defined by

$$\frac{\mathrm{d}w}{\mathrm{d}z} = \frac{\mathrm{d}z}{\mathrm{d}f}, \tag{15}$$

then both e^u and H only depend on the real part of w, thus defining a coupled system of two ODEs with coefficients depending on Q,

$$Q = \frac{2c_\mathrm{q}}{c_\mathrm{h}} \frac{4c_\mathrm{z}\sinh(2c_\mathrm{z}\bar w)}{\sinh(2c_\mathrm{z}w)\sinh(4c_\mathrm{z}\Re(w))},\ \overline{Q} = \frac{2c_\mathrm{q}}{c_\mathrm{h}} \frac{4c_\mathrm{z}\sinh(2c_\mathrm{z}w)}{\sinh(2c_\mathrm{z}\bar w)\sinh(4c_\mathrm{z}\Re(w))}, \tag{16}$$

$$e^u = 4\frac{c_\mathrm{q}}{c_\mathrm{h}^2} v(\Re(w)),\ H = c_\mathrm{h} h(\Re(w)). \tag{17}$$

Therefore the relations (17) define a closed system of ODEs in the independent variable $\xi = \Re(w)$,

$$\begin{cases} (\log v)'' + 8c_\mathrm{q}\left[(h^2 - C)v - \left(\dfrac{4c_\mathrm{z}}{\sinh(4c_\mathrm{z}\xi)}\right)^2 v^{-1}\right] = 0,\ C = \left(\dfrac{c_\mathrm{q}c}{c_\mathrm{h}}\right)^2, \\ vh' + \left(\dfrac{4c_\mathrm{z}}{\sinh(4c_\mathrm{z}\xi)}\right)^2 = 0, \end{cases} \tag{18}$$

which admits a first integral (see Appendix 4 for methods to find first integrals)

$$K = \left(\frac{v'}{v}\right)^2 + 16c_\mathrm{q}\left[(h^2 - C)v + \frac{(4c_\mathrm{z})^2}{\sinh^2(4c_\mathrm{z}\xi)}v^{-1} - 2(4c_\mathrm{z}\coth(4c_\mathrm{z}\xi))h\right] \tag{19}$$

The elimination of v from $(18)_2$ yields a third order first degree ODE [9, §11 p 84 Eq (52)],

$$(\log h')'' + 8c_\mathrm{q}h' - \left(\frac{4c_\mathrm{z}}{\sinh(4c_\mathrm{z}\xi)}\right)^2 \left(8c_\mathrm{q}\frac{h^2 - C}{h'} + 2\right) = 0, \tag{20}$$

and its first integral [12, p. 48] [13, Eq. (4.20)]

$$K = \left(\frac{h''}{h'} + 8c_{\rm z}\coth(4c_{\rm z}\xi)\right)^2 + 16c_{\rm q}\left[\left(\frac{4c_{\rm z}}{\sinh(4c_{\rm z}\xi)}\right)^2\frac{h^2-C}{h'} + h' + 8c_{\rm z}\coth(4c_{\rm z}\xi)h\right]. \tag{21}$$

Therefore, the six arbitrary constants of Bonnet surfaces [9, p. 85] [18, pp. 55] [19] are the two origins of z, $\bar{z}$, the two first integrals $c_{\rm z}$, K, and the two constants of integration of (21).

The integration of this second order second degree ODE could of course not be performed by Bonnet nor Hazzidakis since the complete P6 would only be discovered by R. Fuchs in 1905. This integration, first achieved in [17], is as follows.

When $c_{\rm z} \neq 0$, the ODEs (21) and SD-Ia (A12) obeyed by anyone of the 24 Hamiltonians of P6 (Appendix 4) are equivalent under a homographic transformation, e.g.

$$\begin{cases} h = \dfrac{2c_{\rm z}}{c_{\rm q}}X(X-1)\mathrm{H_{VI}},\ X = \dfrac{1}{1-e^{8c_{\rm z}\xi}}, \\ A_0 = \dfrac{K}{(8c_{\rm z})^2}, A_2 = A_4 = 0,\ A_3 = \dfrac{c_{\rm q}^2 C}{c_{\rm z}^2} - \dfrac{K^2}{4(8c_{\rm z})^2}, \end{cases} \tag{22}$$

in which $\mathrm{H_{VI}}$ is defined in Eq. (A7). The monodromy exponents of P6 are then, for instance (see (A13)),

$$\begin{cases} (\theta_\infty^2, \theta_0^2, \theta_1^2, \Theta_X^2) = (0, t_m + \sqrt{t_m^2 + t_u}, t_m - \sqrt{t_m^2 + t_u}, 0), \\ K = (8c_{\rm z})^2 t_m,\ C = \dfrac{c_{\rm z}^2}{4c_{\rm q}^2}(t_m^2 + t_u). \end{cases} \tag{23}$$

When $c_{\rm z} = 0$, Eq. (21) becomes

$$K = \left(\frac{h''}{h'} + \frac{2}{\xi}\right)^2 + 16c_{\rm q}\left[\frac{h^2-C}{\xi^2 h'} + h' + 2\frac{h}{\xi}\right], \tag{24}$$

and h is then an affine function of the Hamiltonian of P5 [20],

$$-\left(XY''\right)^2 - 4Y'^2(XY'-Y) + A_1(XY'-Y)^2 + A_2(XY'-Y) + A_3Y' + A_4,$$
$$h = \frac{Y_{\rm V}}{4c_{\rm q}\xi},\ \xi = X,\ A_1 = K,\ A_2 = -(8c_{\rm q})^2 C,\ A_3 = 0,\ A_4 = 0. \tag{25}$$

2.2. Constraint to be a surface dual to a Bonnet surface

The involution (11) maps an umbilic-free Bonnet surface (i.e. there exists an isothermic coordinate z such that $1/Q(z,\bar z)$ is harmonic) to another surface which is not a Bonnet surface but which is dual to a Bonnet surface [13, Prop. 4.7.1 page 77] and has a harmonic inverse mean curvature $1/H$ (HIMC).

Let us therefore impose the constraint that $1/H$ be harmonic,

$$(1/H)_{z\bar z}=0. \tag{26}$$

Initially presented in [2, section 7], this constrained system has been integrated with P6 in [10, section 5]. After some algebra, one obtains

$$\frac{1}{H}=\frac{2c_{\rm z}}{c_{\rm h}}\left(\coth(2c_{\rm z}z)+\coth(2c_{\rm z}\bar z)\right)=\frac{2c_{\rm z}}{c_{\rm h}}\frac{\sinh(4c_{\rm z}x)}{\sinh(2c_{\rm z}z)\sinh(2c_{\rm z}\bar z)}, \tag{27}$$

c must be equal to zero, then e^uH^2 and Q are proven to only depend on x [10],

$$\left\{\begin{array}{l} e^u=\dfrac{4c_{\rm u}v(x)}{H^2},\\ Q=\dfrac{4c_{\rm q}}{c_{\rm h}}\left(\dfrac{2c_{\rm z}}{\sinh(2c_{\rm z}x)}\right)^2(q(x)+i\theta),\ \overline{Q}=\dfrac{4c_{\rm q}}{c_{\rm h}}\left(\dfrac{2c_{\rm z}}{\sinh(2c_{\rm z}x)}\right)^2(r(x)-i\theta), \end{array}\right. \tag{28}$$

in which θ is an arbitrary constant. The reduced system [10],

$$\left\{\begin{array}{l} (\log v)''+\dfrac{8}{c_{\rm u}}v-\left(\dfrac{4c_{\rm z}}{\sinh(4c_{\rm z}x)}\right)^2\left(8c_{\rm q}^2\dfrac{(q+i\theta)(r-i\theta)}{c_{\rm u}v}+2\right)=0,\\ c_{\rm u}v-c_{\rm q}q'=0,\\ c_{\rm u}v-c_{\rm q}r'=0, \end{array}\right. \tag{29}$$

admits the first integral

$$\begin{aligned} K=&\left(\frac{v'}{v}+8c_{\rm z}\coth(4c_{\rm z}x)\right)^2\\ &+16\left(\frac{4c_{\rm z}}{\sinh(4c_{\rm z}x)}\right)^2\frac{c_{\rm q}^2}{c_{\rm u}}(q+i\theta)(r-i\theta)+16c_{\rm u}v+16c_{\rm q}\frac{4c_{\rm z}}{\sinh(4c_{\rm z}x)}(q+r). \end{aligned} \tag{30}$$

Without loss of generality, one can choose zero for the constant value of $q-r$. The elimination of v from $(29)_2$ yields a third order first degree ODE for $q(x)$,

$$(\log q')''+8c_{\rm q}q'-\left(\frac{4c_{\rm z}}{\sinh(4c_{\rm z}x)}\right)^2\left(8c_{\rm q}\frac{q^2+\theta^2}{q'}+2\right)=0, \tag{31}$$

which admits the first integral

$$K = \left(\frac{q''}{q'} + 8c_{\mathrm{z}} \coth(4c_{\mathrm{z}}x)\right)^2 + 16c_{\mathrm{q}} \left[\left(\frac{4c_{\mathrm{z}}}{\sinh(4c_{\mathrm{z}}x)}\right)^2 \frac{q^2+\theta^2}{q'} + q' + 8c_{\mathrm{z}} \coth(4c_{\mathrm{z}}x) q\right]. \tag{32}$$

Due to the above-mentioned involution (11), the second order second degree ODEs (21) and (32) are exchanged under

$$h \to q,\ C \to -\theta^2. \tag{33}$$

Therefore (32) integrates with either P6 or P5 depending on whether c_{z} is nonzero or zero, see previous section 2.1.

3. Lie point symmetries and reductions

By requiring the Gauss-Codazzi equations to be invariant under a Lie point symmetry, one can define a reduction to a system (again underdetermined) of three coupled ODEs in four dependent variables. Then, as mentioned in section 2, an additional constraint allows us to integrate this system in terms of P6 functions.

3.1. Lie point symmetries of the Gauss-Codazzi system

Lie point symmetries are obtained by a classical computation presented for instance in [21, 22].

Denoting for convenience $U = e^u, R = \overline{Q}$, the result for the system (9) is the infinite-dimensional Lie algebra generated by

$$\left\{ \begin{array}{l} X(F) = F(z)\partial_z + F'(z)(-2Q\partial_Q - U\partial_U), \\ Y(G) = G(\bar{z})\partial_{\bar{z}} + G'(\bar{z})(-2R\partial_R - U\partial_U), \\ (c = 0 \text{ only})\ a = -H\partial_H + Q\partial_Q + R\partial_R + 2U\partial_U, \end{array} \right. \tag{34}$$

in which F, G are arbitrary functions of one variable. Its table of commutation is

$$\left\{ \begin{array}{l} [X(F_1), X(F_2)] = X(F_1F_2' - F_1'F_2), \\ [Y(G_1), Y(G_2)] = Y(G_1G_2' - G_1'G_2), \\ [X(F), Y(G)] = 0,\ [X(F), a] = 0,\ [Y(G), a] = 0. \end{array} \right. \tag{35}$$

The largest finite-dimensional subalgebra is defined by the seven generators $X(z^j), Y(\bar{z}^j)$, $j = 0, 1, 2$ and a,

$$\left\{ \begin{array}{l} e_0 = \partial_z,\ e_1 = z\partial_z - 2Q\partial_Q - U\partial_U,\ e_2 = z^2\partial_z + 2z(-2Q\partial_Q - U\partial_U), \\ f_0 = \partial_{\bar{z}},\ f_1 = \bar{z}\partial_{\bar{z}} - 2R\partial_R - U\partial_U,\ f_2 = \bar{z}^2\partial_{\bar{z}} + 2\bar{z}(-2R\partial_R - U\partial_U), \\ (c = 0 \text{ only})\ a = -H\partial_H + Q\partial_Q + R\partial_R + 2U\partial_U, \end{array} \right. \tag{36}$$

with the nonzero commutators

$$\begin{array}{l} [e_0, e_1] = e_0,\ [e_0, e_2] = 2e_1,\ [e_1, e_2] = e_2, \\ [f_0, f_1] = f_0,\ [f_0, f_2] = 2f_1,\ [f_1, f_2] = f_2, \end{array} \tag{37}$$

This seven-dimensional algebra is therefore the direct sum

$$\{e_0, e_1, e_2\} \oplus \{f_0, f_1, f_2\} \oplus \{a\}, \tag{38}$$

and the subalgebra $\{e_0, e_1, e_2\}$ is isomorphic to su(1,1), see [23, p. 1451].

There is essentially one resulting reduction, in which the reduced variable is the sum (or, equivalently, the product) of an arbitrary function of z and an arbitrary function of $\bar{z}$.

3.2. Reduction

The infinite-dimensional Lie algebra (34) defines the reduction (with the notation $g_1 = F, g_2 = G$), in which a_1, a_2 denote arbitrary complex constants,

$$\left\{ \begin{array}{l} \xi = \log g_1(z) - \log g_2(\bar{z}),\ e^u = g_1^{2a_1-1} g_2^{2a_2-1} g_1' g_2' \tilde{v},\ H = g_1^{-a_1} g_2^{-a_2} \tilde{h}, \\ Q = g_1^{a_1-2} g_2^{a_2} {g_1'}^2 \tilde{q},\ R = g_2^{a_2-2} g_1^{a_1} {g_2'}^2 \tilde{r}, \end{array} \right. \tag{39}$$

which is indeed a reduction iff the product $a_{\rm s}c$ is zero, with the notation $a_{\rm s} = a_1 + a_2, a_{\rm d} = a_1 - a_2$. The reduced system in $(\tilde{v}, \tilde{h}, \tilde{q}, \tilde{r})(\xi)$,

$$a_{\rm s}c = 0 : \left\{ \begin{array}{l} (\log \tilde{v})'' - \dfrac{1}{2}\left(\tilde{h}^2 - c^2 e^{a_{\rm d}\xi}\right)\tilde{v} + 2\tilde{q}\tilde{r}\tilde{v}^{-1} = 0, \\ 2\tilde{q}' + \tilde{v}\tilde{h}' - \dfrac{1}{2}(a_{\rm s} + a_{\rm d})\tilde{v}\tilde{h} - (a_{\rm s} - a_{\rm d})q = 0, \\ 2\tilde{r}' + \tilde{v}\tilde{h}' + \dfrac{1}{2}(a_{\rm s} - a_{\rm d})\tilde{v}\tilde{h} + (a_{\rm s} + a_{\rm d})r = 0, \end{array} \right. \tag{40}$$

whose reduced moving frame equations are given in the Appendix section 4, admits the first integral

$$K = \left(\frac{\tilde{v}'}{\tilde{v}} + a_{\rm d}\right)^2 - \left(h^2 - c^2 e^{a_{\rm d}\xi}\right)\tilde{v} - 2\tilde{h}(\tilde{q} + \tilde{r}) - 2ce^{a_{\rm d}\xi/2}(\tilde{q} - \tilde{r}) - 4\frac{\tilde{q}\tilde{r}}{\tilde{v}}\cdot \tag{41}$$

Except when $a_{\rm d}c \neq 0$, the system (40) and its first integral are autonomous.

Under the change of variables defined by

$$e^{\xi} = \eta,\ \tilde{v} = e^{(2a_2-1)\xi} v,\ \tilde{h} = e^{-a_2\xi} h,\ \tilde{q} = e^{a_2\xi} q,\ \tilde{r} = e^{(a_2-2)\xi} r,\ a_{\rm s} = g + 1, \tag{42}$$

the system (40)–(41) is mapped to a system with one less parameter,

$$(1+g)c = 0 : \left\{ \begin{array}{l} \left(\eta\dfrac{v'}{v}\right)' - \dfrac{h^2 - c^2}{2\eta^2} v + 2\dfrac{qr}{\eta^2 v} = 0, \\ -\eta v h' - 2\eta^2 q' + (1+g) v h = 0, \\ -\eta^2 v h' - 2\eta r' + 2(1-g) r = 0, \end{array} \right. \tag{43}$$

which admits the first integral

$$K = \left(\eta\frac{v'}{v} + g\right)^2 - \frac{(h^2 - c^2)v}{\eta} - 2(h + c)q - 2\frac{(h-c)}{\eta^2} r - 4\frac{qr}{\eta v}\cdot \tag{44}$$

From now on, we impose the constraint $q = r$ (equivalent to the isothermic constraint $Q = \overline{Q} = R$ under a suitable choice of the so far arbitrary functions $g_1(z), g_2(\bar{z})$).

3.2.1. Reduction, generic isothermic case, $g^2 \neq 1, c = 0$. In the generic case $g^2 \neq 1$, c must vanish and the three fields $hv/q, h'/h, q'/q$ are equivalent under the homographic group,

$$g^2 \neq 1 : \ 1 + \frac{\eta h v}{2q} = \frac{2 - (1-\eta^2)\eta h'/h}{(1+g)} = 1 - \frac{(1-g)\eta^2}{(1-\eta^2)\eta q'/q - (1-g)}, \quad (45)$$

therefore it is more convenient to choose them as dependent variables rather than v, h, q. After elimination of two of these three variables $hv/q, h'/h, q'/q$, the remaining variable is found to obey a second order first degree ODE. Since the three such ODEs only differ by the homographic transformation (45), it is sufficient to consider only one of them. In the case $W = \eta h v/(2q)$ for instance, this is,

$$\frac{\mathrm{d}^2 W}{\mathrm{d}\eta^2} = \frac{1}{2}\left[\frac{1}{W} + \frac{1}{W+1} + \frac{1}{W+\eta^2}\right]\left(\frac{\mathrm{d}W}{\mathrm{d}\eta}\right)^2$$
$$- \left[\frac{1}{\eta} - \frac{2\eta}{1-\eta^2} - \frac{2\eta}{W+\eta^2}\right]\frac{\mathrm{d}W}{\mathrm{d}\eta} + \frac{2W(W+1)(W+\eta^2)}{\eta^2(1-\eta^2)^2} \times$$
$$\left[\left(\frac{1+g}{2}\right)^2 - \left(\frac{1-g}{2}\right)^2 \frac{\eta^2}{4W^2} + \frac{K}{4}\frac{\eta^2-1}{(W+1)^2} + (1-\frac{K}{4})\frac{\eta^2(\eta^2-1)}{(W+\eta^2)^2}\right]. \quad (46)$$

This ODE belongs to the class studied by Painlevé and Gambier [24]

$$W'' = A_2(W,\eta)W'^2 + A_1(W,\eta)W' + A_0(W,\eta), \quad (47)$$

the rational function A_2 of W is the sum of four simple poles with equal residues $1/2$, located at

$$W = \infty, \ 0, \ -1, \ -\eta^2, \quad (48)$$

these four poles are distinct and their crossratio is not a constant, therefore the ODE (46) is a homographic transform of P6, Eq. (A6). This provides the general solution in terms of a P6 function $V(X)$ with two constraints among its four parameters θ_j, i.e. as many as in the two ex-

isting solutions,

$$\left\{\begin{array}{l} W=\dfrac{\eta h v}{2q}=\dfrac{V-X}{X},\\ \dfrac{\mathrm{d}\log h}{\mathrm{d}X}=\dfrac{1+g}{2(X-1)}-\dfrac{1-g}{2(V-X)},\ \dfrac{\mathrm{d}\log q}{\mathrm{d}X}=\dfrac{1}{X-1}-\dfrac{(1+g)V}{2X(X-1)},\\ hq=\dfrac{2(X-1)}{V(V-1)(V-X)}\\ \qquad\times\left[\left(X(X-1)V'+\dfrac{1-g}{2}V(V-1)\right)^2-\dfrac{K}{4}(V-X)^2\right],\\ X=\dfrac{1}{1-\eta^2},\ (\theta_\infty^2,\theta_0^2,\theta_1^2,\theta_X^2)=\left(\left(\dfrac{1+g}{2}\right)^2,\dfrac{K}{4},\dfrac{K}{4},\left(\dfrac{1-g}{2}\right)^2\right). \end{array}\right. \tag{49}$$

Several remarks are in order.

1. This solution is invariant under the involution (11), which acts as a homography on V.
2. For generic values of the θ_j's, there exists no rational function of V, V' which would represent h or q. Only their product hq is rational.
3. Since there exists a choice of square roots allowing the monodromy exponents θ_j to have an integral sum, there always exists a one-parameter solution of this P6 which is an algebraic transform of the hypergeometric function and obeys, for instance,
$$g^2\neq 1:\ \left\{\begin{array}{l} \dfrac{X(X-1)V'}{V(V-1)(V-x)}+\dfrac{\theta_0}{V}+\dfrac{\theta_1}{V-1}+\dfrac{\theta_X-1}{V-X}=0,\\ \theta_\infty=\dfrac{1+g}{2},\ \theta_0=\dfrac{\sqrt{K}}{2},\ \theta_1=-\dfrac{\sqrt{K}}{2},\ \theta_X=\dfrac{1-g}{2}. \end{array}\right. \tag{50}$$
4. The values of the monodromy exponents even allow algebraic solutions to exist. For instance, the cube solution of P6 [25]
$$\left\{\begin{array}{l} -X^2+3XV-3XV^2+2XV^3-V^3=0\\ (\theta_\infty,\theta_0,\theta_1,\theta_X)=(1/3,a,a,2a),\ a \text{ arbitrary}, \end{array}\right. \tag{51}$$
matches (49) for $a=2/3$, $g=-1/3$, $K=16/9$.
5. From the definitions (39) and (42), we could not build a function of (e^u,H,Q) which would be a harmonic function of $(z,\bar z)$, like for instance $z^2(\eta^2-1)/\eta^2$.
6. One may wonder whether the invariances of P6 allow mapping the present surface to a Bonnet surface. The answer is negative, at least for generic values of the monodromy exponents. Indeed, the elementary birational transformation which conserves P6 [26] acts as an affine transformation on the θ_j's (see, e.g. [27, §B.3.1]) and, starting from the θ_j's in (49), one can easily make one transformed θ_j equal to zero, but not two.

3.2.2. Reduction, isothermic case, $g=1, c=0$. The elimination of v among the last two equations of (43) defines two subcases,

$$q' \left[(1-\eta^2)\eta h' - 2h \right] = 0. \tag{52}$$

The first subcase $q'=0$ is equivalent to

$$g=1,\ c=0,\ h=0,\ q'=0,\ \left(\eta\frac{v'}{v}+1\right)^2 - K - 4\frac{q^2}{\eta v} = 0, \tag{53}$$

and its general solution

$$g=1,\ c=0,\ h=0,\ q=c_{\rm q},\ \eta=e^{\xi},\ \eta v = \begin{cases} \dfrac{4c_{\rm q}^2}{K}\sinh^2(\dfrac{\sqrt{K}}{2}(\xi-\xi_0)),\ c_{\rm q}K\neq 0, \\ c_{\rm q}^2(\xi-\xi_0)^2,\ c_{\rm q}\neq 0,\ K=0, \\ v_0 e^{\sqrt{K}(\xi-\xi_0)},\ c_{\rm q}=0, \end{cases} \tag{54}$$

depends on at most three movable constants among $K, c_{\rm q}, \xi_0, v_0$.

The second subcase $q'\neq 0$ is equivalent to

$$g=1,\ c=0,\ h=c_{\rm h}\frac{\eta^2}{\eta^2-1},\ v=-\frac{q'}{h},\ c_{\rm h}\neq 0, \tag{55}$$

in which q is the Hamiltonian of P6, see Eqs. (A10)–(A11),

$$q(\eta) = -\frac{8}{c_{\rm h}}X(X-1)\mathrm{H_{VI}},\ X=\frac{1}{1-\eta^2},\ A_0=\frac{K}{4},\ A_2=0,\ A_3=0,\ A_4=0, \tag{56}$$

and the four integration constants are $K, c_{\rm h}$ and the two movable constants of P6. This solution, for which $1/H$ is harmonic,

$$\frac{1}{H} = \frac{1}{c_{\rm h}}\left(g_1^2(z) - g_2^2(\bar z)\right), \tag{57}$$

is the particular case $\theta=0$ of the HIMC solution of section 2.2.

3.2.3. Reduction, isothermic case, $g=-1, c$ arbitrary. The elimination of v among the last two equations of (43) defines two subcases,

$$h' \left[(1-\eta^2)\eta q' - 2q \right] = 0. \tag{58}$$

The first subcase $h'=0$ is equivalent to

$$g=-1,\ c \text{ arbitrary},\ h'=0,\ q=0,\ \left(\eta\frac{v'}{v}-1\right)^2 - K - (h^2-c^2)\frac{v}{\eta} = 0, \tag{59}$$

and its general solution

$$g=-1,\ h=h_0,\ q=0,\ \frac{\eta}{v}=\begin{cases}\dfrac{h_0^2-c^2}{K}\sinh^2((\sqrt{K}/2)\xi-\xi_0),\ (h_0^2-c^2)K\neq 0,\\ \dfrac{h_0^2-c^2}{4}(\xi-\xi_0)^2,\ (h_0^2-c^2)\neq 0,\ K=0,\\ v_0e^{\sqrt{K}(\xi-\xi_0)},\ h_0^2-c^2=0,\end{cases}\tag{60}$$

depends on at most three movable constants among K, h_0, ξ_0, v_0.

The second subcase $h'\neq 0$ is equivalent to

$$g=-1,\ c\text{ arbitrary},\ q=c_{\rm q}\frac{\eta^2}{\eta^2-1},\ v=\frac{4c_{\rm q}\eta^2}{(1-\eta^2)^2h'},\ c_{\rm q}\neq 0,\tag{61}$$

in which h is an affine function of the Hamiltonian of P6, see Eqs. (A10)–(A11),

$$\begin{cases}h=-\dfrac{8}{c_{\rm q}}X(X-1)\mathrm{H_{VI}},\ X=\dfrac{1}{1-\eta^2},\\ A_0=\dfrac{K+2c_{\rm q}c}{4},\ A_2=0,\ A_3=-\dfrac{K(K+4c_{\rm q}c)}{64}A_4=0,\end{cases}\tag{62}$$

and the four integration constants are $K, c_{\rm q}$ and the two movable constants of P6. This solution, in which $1/Q$ and $1/R$ are harmonic up to some conformal transformation,

$$\frac{(g_1'(z))^2}{Q}=\frac{(g_2'(\bar z))^2}{R}=\frac{1}{c_{\rm q}}\left(g_1^2(z)-g_2^2(\bar z)\right),\tag{63}$$

is identical to the solution of Bonnet described in section 2.1.

These two nongeneric solutions $g=1,-1$ are of course exchanged under the involution (11).

In Table 1, we have gathered all the (complex) solutions of the $\mathbb{R}^3(c)$ Gauss-Codazzi equations which involve P6 or P5. Let us remind once again that all constants $c_{\rm z}, c_{\rm u}, c_{\rm q}$, etc, are complex. For instance, $4c_{\rm z}/\sinh(4c_{\rm z}x)$ denotes at the same time $1/\sinh x$, $1/\sin x$ or $1/x$.

Table 1

Complex solutions in terms of Painlevé P6 or P5 functions.
This table collects both old and new such solutions. Other solutions (in terms of classical functions) can be found in the text. The second column displays the function which is harmonic. The column "Link" refers to the explicit link to a Painlevé function given in the text. The column "arb" lists the arbitrary constants in P6 or P5.

Reduction	Harmo	e^u	H	Q	Integration	c	arb	Link	Ref
x	$1/Q$	$\left(\frac{4c_z}{\sinh(4c_z x)}\right)^2 h'(x)^{-1}$	$h(x)$	$\frac{4c_z \sinh(2c_z \bar{z})}{\sinh(2c_z z)\sinh(4c_z x)}$	$h(x) = \mathrm{H_{VI}}$	c	K, c	(22)	[9]
x	$1/Q$	$x^{-2} h'(x)^{-1}$	$h(x)$	$\bar{z}/(zx)$	$h(x) = \mathrm{H_V}$	c	K, c	(25)	[9]
x	$1/H$	$q'(x) H^{-2}$	$c_\mathrm{h} \frac{\sinh(2c_z \bar{z})\sinh(2c_z z)}{2c_z \sinh(4c_z x)}$	$\left(\frac{2c_z}{\sinh(2c_z x)}\right)^2 (q + i\theta)$	$q(x) = \mathrm{H_{VI}}$	0	K, θ	(22)	[10]
x	$1/H$	$q'(x) H^{-2}$	$z\bar{z}/(2x)$	$x^{-2}(q(x) + i\theta)$	$q(x) = \mathrm{H_V}$	0	K, θ	(25)	[10]
$\eta = z/\bar{z}$	?	$z^{2g} v(\eta)$	$z^{-1-g} h(\eta)$	$z^{-1+g} q(\eta)$	Reduction (43)	0			
$g = 1$	$1/H$	$-z^2 q'(\eta) \frac{\eta^2 - 1}{\eta^2}$	$z^{-2}\eta^2/(\eta^2 - 1)$	$q(\eta)$	$q(\eta) = \mathrm{H_{VI}}$	0	K	(56)	[10]
$g = -1$	$1/Q$	$z^{-2} \frac{\eta^2}{(1-\eta^2)^2 h'(\eta)}$	$h(\eta)$	$z^{-2} \frac{\eta^2}{\eta^2 - 1}$	$h(\eta) = \mathrm{H_{VI}}$	c	K, c	(62)	[9]
$g^2 \neq 1$	?	$z^{2g} v(\eta)$	$z^{-1-g} h(\eta)$	$z^{-1+g} q(\eta)$	$h'/h, q'/q, hv/q = \mathrm{P6}$	0	K, g	(49)	this paper

3.3. On the linear representations of P6

In each of the three cases $g^2 \neq 1$, $g = 1$, $g = -1$, the reduced moving frame equations (A5) define a linear representation of a codimension-two P6. For $g^2 = 1$, these have been shown [13, §3.4.1 and 4.9.1] to be essentially not different from the isomonodromic matrix Lax pairs of Jimbo and Miwa, the spectral parameter t being defined by (A4).

In the case $g^2 \neq 1$, the reduced moving frame equations (A5) do not depend on the parameter t and therefore define a linear representation of P6 which is not a Lax pair. This traceless, well-balanced, representation is,

$$
\begin{aligned}
&\mathrm{d}\Psi = \tilde{U}_r\Psi\mathrm{d}\xi + \tilde{V}_r\Psi\mathrm{d}t = U_r\Psi\mathrm{d}X + V_r\Psi\mathrm{d}t,\\
&U_r = \frac{1}{4X(X-1)}\left[\frac{1+g}{2}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\right.\\
&\qquad\left.+\frac{\sqrt{R_+R_-}}{\sqrt{V(V-1)}(V-X)}\begin{pmatrix}0 & V-2X+1\\ V-2X & 0\end{pmatrix}\right],\\
&V_r = \frac{X(X-1)V' - V(V-1)(1+g)/2}{2(V-X)}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}\\
&\qquad+\frac{\sqrt{R_+R_-}}{2\sqrt{V(V-1)}(V-X)}\begin{pmatrix}0 & -(V-1)\\ V & 0\end{pmatrix},\\
&R_\pm = X(X-1)\frac{\mathrm{d}V}{\mathrm{d}X} + V(V-1)(V-X)\left[\pm\frac{\theta_0}{V} \mp \frac{\theta_1}{V-1} + \frac{\theta_X - 1}{V-X}\right],\\
&(\theta_\infty^2, \theta_0^2, \theta_1^2, \theta_X^2) = \left(\left(\frac{1+g}{2}\right)^2, \frac{K}{4}, \frac{K}{4}, \left(\frac{1-g}{2}\right)^2\right),
\end{aligned}
\tag{64}
$$

and it is not too difficult to extend it to the generic P6.

One such extrapolation

$$
\begin{aligned}
U_r &= \frac{a_{11}}{4}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix} + \frac{(V-2X+1)\sqrt{R}}{4X(X-1)\sqrt{V(V-1)}(V-X)}\begin{pmatrix}0 & 1\\ 0 & 0\end{pmatrix}\\
&\quad + \frac{a_{21}}{4X(X-1)\sqrt{RV(V-1)}}\begin{pmatrix}0 & 0\\ 1 & 0\end{pmatrix},\\
a_{11} &= -\left(\frac{1}{V}+\frac{1}{V-1}+\frac{1}{V-X}\right)\frac{\mathrm{d}V}{\mathrm{d}X} + \frac{\mathrm{d}\log R}{\mathrm{d}X} + \frac{2}{V-X}\\
&\quad + \left(\frac{-2}{V-1}+\frac{1}{V-X}\right)F,\\
a_{21} &= \frac{V-2X}{V-X}\left[X(X-1)\frac{\mathrm{d}V}{\mathrm{d}X}\right]^2\\
&\quad + 2X(X-1)\frac{\mathrm{d}V}{\mathrm{d}X}\left[2V\partial_V F - \left(\frac{2}{V-1}+\frac{1}{V-X}\right)F\right]\\
&\quad + 4X(X-1)V\left(\partial_X F + \frac{F}{V-X}\right) + 2X(X-1)\frac{V^2(V-1)}{V-X}\\
&\quad + V\left(-\frac{2}{V-1}+\frac{1}{V-X}\right)\left(F^2+k(V-X)^2\right)\\
&\quad + 2V^2(V-1)(V-X)\left[\theta_\infty^2 - \theta_0^2\frac{X}{V^2} + \theta_1^2\frac{X-1}{(V-1)^2} - \theta_X^2\frac{X(X-1)}{(V-X)^2}\right],\\
V_r &= \frac{X(X-1)V'+F}{2(V-X)}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix} - \frac{(V-1)\sqrt{R}}{2\sqrt{V(V-1)}(V-X)}\begin{pmatrix}0 & 1\\ 0 & 0\end{pmatrix}\\
&\quad + \frac{V\left[(X(X-1)V'+F)^2 + k(V-X)^2\right]}{2\sqrt{RV(V-1)}(V-1)}\begin{pmatrix}0 & 1\\ 0 & 0\end{pmatrix},
\end{aligned}
\tag{65}
$$

depends on two arbitrary functions R and $F(V,X)$ and one arbitrary constant k, and it reduces to (64) for the values

$$
R = R_+R_-,\ F = -\frac{1+g}{2}V(V-1),\ k = -\frac{K}{4}\cdot \tag{66}
$$

4. Conclusion

There exist at least two types of solutions of the Gauss-Codazzi system (9) in terms of P6. In the first type, the set $\{1/H, 1/Q\}$ is made of a harmonic function and the Hamiltonian of P6. These are the solutions of Bonnet (1867) and Bobenko, Eitner and Kitaev (1997). The solution $1/Q$ harmonic defines a Bonnet surface, while the solution $1/H$ harmonic defines a surface dual to a Bonnet surface [13, p. 77].

The second type is presented for the first time in this article. In this type, the product $e^u H/Q$ and the logarithmic derivatives of H and Q with respect to the reduced independent variable are homographic transforms of the same P6 function, and there is apparently no simple geometric variable which is harmonic. The only such solution which we could find is (49), and it requires two constraints among the four parameters of P6, i.e. the same number as the Bonnet and BEK solutions. The corresponding analytic surface is isothermic but is neither a Bonnet surface nor a surface dual to a Bonnet surface.

Another point worth being emphasized is the power of the Lie approach. Indeed, among the set of infinitesimal point symmetries, there exists at least one algebra able to recover the two previously known P6 solutions (Bonnet 1867, Bobenko, Eitner and Kitaev 1997), see (56), (62). However, a limitation of this Lie approach is its failure to introduce a spectral parameter in the reduced Gauss-Weingarten equations (64).

In future research, we plan to remove the two constraints among the four parameters of P6.

Acknowledgments

RC warmly acknowledges the generous support of both the Centre de recherches mathématiques and l'Unité mixte internationale 3457 (CRM) du CNRS. RC was also partially supported by the Laboratoire de recherche conventionné LRC-Méso. AMG is pleased to thank Digiteo for financial support. We thank Hsieh Chun-Chung and Ma Hui for stimulating discussions.

Appendix A. First integrals and reduced moving frames

There are two methods to find a first integral to a third order first degree ODE such as (20), (31).

The first one relies on classical results [11]. If such a third order ODE admits an algebraic first integral and if the second order ODE defined by this first integral has the Painlevé property, then its degree d (in the second derivative) is bounded and can only take the values $d = 1, 2, 3, 4, 6$ of Briot and Bouquet. The first integral then has the necessary form

$$K = \sum_{j=0}^{d} F_j(h', h, x) \left(h''\right)^j , \tag{A1}$$

which makes it easy to compute. In all cases handled in this paper, the degree is $d = 2$. This is probably the method used by Hazzidakis to obtain the first integral of (20).

The second method is to take advantage of the linear representation by the moving frame (Gauss-Weingarten) equations. Let us take the system (40) as an example. Under the reduction (39), the PDE moving frame equations (7)

$$\mathrm{d}\psi = U\psi \mathrm{d}z + V\psi \mathrm{d}\bar{z} \tag{A2}$$

is mapped to an ODE moving frame

$$\mathrm{d}\Psi = \tilde{U}_r \Psi \mathrm{d}\xi + \tilde{V}_r \Psi \mathrm{d}t, \tag{A3}$$

in which the parameter t and the transition matrix P are defined as (see [27, §5.8.1] for the systematic derivation of t and P),

$$\begin{aligned} &t = \log g_1(z) + \log g_2(\bar{z}),\ \psi = P\Psi, \\ &P = \mathrm{diag}(((\log g_1)')^{1/4}((\log g_2)')^{-1/4}, ((\log g_1)')^{-1/4}((\log g_2)')^{1/4}). \end{aligned} \tag{A4}$$

These ODE moving frame equations

$$\begin{aligned} \tilde{U}_r &= \frac{1}{2}\begin{pmatrix} a_\mathrm{s}/2 & \left((\tilde{h} - ce^{a_\mathrm{d}\xi/2})/2 - \tilde{q}/\tilde{v}\right)\tilde{v}^{1/2} \\ \left((\tilde{h} + ce^{a_\mathrm{d}\xi/2})/2 - \tilde{r}/\tilde{v}\right)\tilde{v}^{1/2} & -a_\mathrm{s}/2 \end{pmatrix}, \\ \tilde{V}_r &= \frac{1}{2}\begin{pmatrix} a_\mathrm{d}/2 + \tilde{v}'/(2\tilde{v}) & \left(-(\tilde{h} - ce^{a_\mathrm{d}\xi/2}) - \tilde{q}/\tilde{v}\right)\tilde{v}^{1/2} \\ \left((\tilde{h} + ce^{a_\mathrm{d}\xi/2}) + \tilde{r}/\tilde{v}\right)\tilde{v}^{1/2} & -a_\mathrm{d}/2 - \tilde{v}'/(2\tilde{v}) \end{pmatrix} \end{aligned} \tag{A5}$$

then generate the first integral $K = 8\,\mathrm{tr}\,\tilde{V}_r^2$, Eq. (41).

Appendix B. Hamiltonian and τ-function of P6

The P6 function is by definition the general solution of the P6 equation,

$$\frac{\mathrm{d}^2V}{\mathrm{d}X^2} = \frac{1}{2}\left[\frac{1}{V} + \frac{1}{V-1} + \frac{1}{V-X}\right] V'^2 - \left[\frac{1}{X} + \frac{1}{X-1} + \frac{1}{V-X}\right] V' + \frac{V(V-1)(V-X)}{2X^2(X-1)^2}\left[\theta_\infty^2 - \theta_0^2\frac{X}{V^2} + \theta_1^2\frac{X-1}{(V-1)^2} + (1-\theta_X^2)\frac{X(X-1)}{(V-X)^2}\right], \tag{A6}$$

which contains four equivalent singularities $V = \infty, 0, 1, X$. Let us recall the main property of P6. This is the *unique* function defined by the class of second order algebraic ordinary differential equations (ODE). Indeed, all such ODEs which possess the Painlevé property have for general solution either an algebraic transform of P6 or of one of its five degeneracies P5, ..., P1, or an elliptic function, or the solution of some linear ODE.

For each $\mathrm{Pn}(V, X)$, there exists a rational function $\mathrm{H}(V, V', X)$ whose only movable singularities are simple poles of residue unity, thus defining a function $\tau = e^{\int \mathrm{H}\,\mathrm{d}X}$, called *$\tau$-function*, which has the same kind of movable singularities as the general solution of a linear ODE (namely no movable singularities at all, only movable simple zeroes).

For P6, there exist 24 choices for H. Rather than the expression built by Painlevé [28, Eq. (3)] and Chazy [29, t page 341], it is preferable to adopt the choice of Malmquist [30],

$$\mathrm{H}_{\mathrm{VI,M}} = \frac{X(X-1)V'^2}{4V(V-1)(V-X)} + \frac{1}{4X(X-1)}\left[-\theta_\infty^2\left(V - \frac{1}{2}\right) - \theta_0^2\left(\frac{X}{V} - \frac{1}{2}\right) + \theta_1^2\left(\frac{X-1}{V-1} - \frac{1}{2}\right) - (\theta_X - 1)^2\left(\frac{X(X-1)}{V-X} + X - \frac{1}{2}\right)\right]. \tag{A7}$$

The conjugate variables q, p

$$q = V,\ 2p = \frac{X(X-1)V'}{V(V-1)(V-X)} + \frac{\theta_0}{V} + \frac{\theta_1}{V-1} + \frac{\theta_X - 1}{V-X}, \tag{A8}$$

then define the polynomial Hamiltonian $H(q, p, x) = \mathrm{H}(u, u', x)$ [30]

$$H = \frac{q(q-1)(q-X)}{X(X-1)}\left[p^2 - \left(\frac{\theta_0}{q} + \frac{\theta_1}{q-1} + \frac{\theta_X - 1}{q-X}\right)p - \frac{\theta_\infty^2 - (\theta_0 + \theta_1 + \theta_X - 1)^2}{4q(q-1)}\right]. \tag{A9}$$

The 24 $\mathrm{H_{VI}}$ obey the same second order second degree ODE, labeled (B-V) by Chazy [29, page 340]. After the normalization

$$Y = X(X-1)\mathrm{H_{VI}}, \tag{A10}$$

the ODE for $Y(X)$ can be written either with a nice quaternary symmetry [31],

$$\begin{aligned} -Y'\left(X(X-1)Y''\right)^2 - \left[Y'^2 - 2Y'(XY'-Y) + n_1n_2n_3n_4\right]^2 \\ + (Y'+n_1^2)(Y'+n_2^2)(Y'+n_3^2)(Y'+n_4^2) = 0, \end{aligned} \tag{A11}$$

or as a simplified equation with four complementary terms [20, Eq. (5.4)],

$$\begin{aligned} \text{SD-Ia}: \ -\left((X(X-1)Y''\right)^2 - 4Y'(XY'-Y)^2 + 4Y'^2(XY'-Y) \\ + A_0Y'^2 + A_2(XY'-Y) + \left(A_3 + \frac{A_0^2}{4}\right)Y' + A_4 = 0, \end{aligned} \tag{A12}$$

with the correspondence,

$$\left\{ \begin{array}{l} \Theta_X = \theta_X - 1, \\ 2n_1 = \theta_\infty - \Theta_X,\ 2n_2 = \theta_\infty + \Theta_X,\ 2n_3 = \theta_1 - \theta_0,\ 2n_4 = \theta_1 + \theta_0, \\ 2A_0 = \theta_\infty^2 + \theta_0^2 + \theta_1^2 + \Theta_X^2 = 2(n_1^2 + n_2^2 + n_3^2 + n_4^2), \\ 4A_2 = -(\theta_\infty^2 - \Theta_X^2)(\theta_0^2 - \theta_1^2) = 16n_1n_2n_3n_4, \\ 4A_3 = (\theta_\infty^2 - \theta_1^2)(\theta_0^2 - \Theta_X^2) \\ \qquad = -(n_1+n_2+n_3+n_4)(n_1+n_2-n_3-n_4) \\ \qquad\quad \times (n_1-n_2+n_3-n_4)(n_1-n_2-n_3+n_4), \\ 32A_4 = (\theta_\infty^2 + \Theta_X^2)(\theta_0^2 - \theta_1^2)^2 + (\theta_\infty^2 - \Theta_X^2)^2(\theta_0^2 + \theta_1^2) \\ \qquad = 32(n_1n_2n_3n_4)^2(n_1^{-2} + n_2^{-2} + n_3^{-2} + n_4^{-2}). \end{array} \right. \tag{A13}$$

For reference, the inverse transformation is [31, Table R],

$$\begin{aligned} V = X + \frac{\Theta_X X(X-1)Y''}{(Y' + (\theta_\infty + \Theta_X)^2/4)(Y' + (\theta_\infty - \Theta_X)^2/4)} \\ + \frac{r_+}{Y' + (\theta_\infty + \Theta_X)^2/4} + \frac{r_-}{Y' + (\theta_\infty - \Theta_X)^2/4}, \\ r_+ + r_- = -Y - \frac{\theta_\infty^2 + 3\Theta_X^2}{8}(2X-1) + \frac{\theta_1^2 - \theta_0^2}{8}, \\ r_+ + r_- = \left(-Y - \frac{3\theta_\infty^2 + \Theta_X^2}{8}(2X-1) - \frac{\theta_1^2 - \theta_0^2}{8}\right)\frac{\Theta_X}{\theta_\infty}\cdot \end{aligned} \tag{A14}$$

References

1. L. Bianchi, Sulle superficie a linee di curvatura isoterme, Rend. Acc. Naz. dei Lincei **12** (1903) 511–520.
2. A.I. Bobenko, Surfaces in terms of 2 by 2 matrices: Old and new integrable cases, 83–128, in *Harmonic maps and integrable systems*, A.P. Fordy and J.C. Wood (eds.), Aspects of mathematics E23, Vieweg, Braunschweig, Wiesbaden, 1994.
3. Boris A. Springborn, Bonnet pairs in the 3-sphere, Contemporary mathematics **308** (2002) 297–303.
4. A. Sym, Soliton surfaces, Lett. Nuovo Cimento **33** (1982) 394–400.
5. A. Sym, Soliton surfaces and their applications, 154–231, *Geometrical aspects of the Einstein equations and integrable systems*, Lecture notes in physics **239**, ed. R. Martini (Springer, Berlin, 1985).
6. J.L. Cieśliński, P. Goldstein and A. Sym, Isothermic surfaces in E^3 as soliton surfaces, Phys. Lett. A **205** (1995) 37–43.
7. A.I. Bobenko, Exploring surfaces through methods from the theory of integrable systems. Lectures on the Bonnet problem, School on differential geometry, 12–30 April 1999, Trieste.
   ftp://ftp-sfb288.math.tu-berlin.de/pub/Preprints/preprint403.ps.gz
8. Chunli Chen and Yishen Li, The Lund-Regge surface and its motion's evolution equation, J. Math. Phys. **43** (2002) 1938–1955.
9. O. Bonnet, Mémoire sur la théorie des surfaces applicables sur une surface donnée. Deuxième partie : Détermination de toutes les surfaces applicables sur une surface donnée, J. École polytechnique **42** (1867) 1–151.
   http://gallica.bnf.fr/ark:/12148/bpt6k433698b/f5.image
10. A.I. Bobenko, U. Eitner and A.V. Kitaev, Harmonic inverse mean curvature surfaces and Painlevé equations, Geometriae dedicata **68** (1997) 187–227.
11. P. Painlevé, Mémoire sur les équations différentielles dont l'intégrale générale est uniforme, Bull. Soc. Math. France **28** (1900) 201–261.
12. J.N. Hazzidakis, Biegung mit Erhaltung der Hauptkrümmungsradien, Journal fï die reine und angewandte Mathematik **117** (1897) 42–56.
13. A.I. Bobenko and U. Eitner, Painlevé equations in differential geometry of surfaces, 120 pages, Lecture Notes in Math. **1753** (Springer, Berlin, 2000).
14. Weihuan Chen and Haizhong Li, Bonnet surfaces and isothermic surfaces, Result. Math. **31** (1997) 40–52.
15. W.C. Graustein, Applicability with preservation of both curvatures, Bull. Amer. math. soc. **30** (1924) 19–23.
16. R. Tribuzy, A characterization of tori with constant mean curvature in a space form, Bol. Soc. Brasil. Mat. **11** (1980) 259–274.
17. A.I. Bobenko and U. Eitner, Bonnet surfaces and Painlevé equations, Journal für die reine und angewandte Mathematik **499** (1998) 47–79.
18. É. Cartan, Sur les couples de surfaces applicables avec conservation des courbures principales, Bulletin des sciences mathématiques **66** (1942) 55–72, 74–85.
19. S.S. Chern, Deformation of surfaces preserving principal curvatures, *Differential geometry and complex analysis*, 155–163, eds. I. Chavel and H.M. Farkas, H.E. Rauch memorial volume (Springer, Berlin, 1985).
20. C.M. Cosgrove and G. Scoufis, Painlevé classification of a class of differential equations of the second order and second degree, Stud. Appl. Math. **88** (1993) 25–87.

21. P.J. Olver, *Applications of Lie groups to differential equations* (Springer, Berlin, 1986).
22. L.V. Ovsiannikov, *Group properties of differential equations*, (Siberian section of the Academy of Sciences of the USSR, Novosibirsk, 1962) in Russian. Translated by G.W. Bluman (1967), *Group analysis of differential equations* (Academic press, New York, 1982).
23. J. Patera and P. Winternitz, Subalgebras of real three- and four-dimensional Lie algebras, J. Math. Phys. **18** (1977) 1449–1455.
24. B. Gambier, Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes, Acta Math. **33** (1910) 1–55.
25. A.V. Kitaev, Special functions of the isomonodromy type, rational transformations of spectral parameter, and algebraic solutions of the sixth Painlevé equation, Algebra i Analiz **14** (2002) 121–139. English translation: St. Petersburg Math. J. **14** (2003) 453–465. http://arXiv.org/abs/nlin.SI/0102020
26. K. Okamoto, Studies on the Painlevé equations, I, Sixth Painlevé equation, Ann. Mat. Pura Appl. **146** (1987) 337–381.
27. R. Conte and M. Musette, *The Painlevé handbook* (Springer, Berlin, 2008). Russian translation *Metod Penleve y ego prilozhenia* (Regular and chaotic dynamics, Moscow, 2011).
28. P. Painlevé, Sur les équations différentielles du second ordre à points critiques fixes, C. R. Acad. Sc. Paris **143** (1906) 1111–1117.
29. J. Chazy, Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'intégrale générale a ses points critiques fixes, Acta Math. **34** (1911) 317–385.
30. J. Malmquist, Sur les équations différentielles du second ordre dont l'intégrale générale a ses points critiques fixes, Arkiv för Math. Astr. Fys. **17** (1922–23) 1–89.
31. K. Okamoto, Polynomial Hamiltonians associated with Painlevé equations. II, Differential equations satisfied by polynomial Hamiltonians, Proc. Japan Acad. A **56** (1980) 367–371.

1. Centre de mathématiques et de leurs applications
École normale supérieure de Cachan, CNRS, Université Paris-Saclay,
61, avenue du Président Wilson, F–94235 Cachan Cedex, France.

2. Department of Mathematics, The University of Hong Kong,
Pokfulam Road, Hong Kong.

3. Centre de recherches mathématiques, Université de Montréal
Case postale 6128, Succursale Centre ville,
Montréal, Québec H3C 3J7, Canada

4. Département de mathématiques et d'informatique,
Université du Québec à Trois-Rivières
Case postale 500, Trois-Rivières, Québec G9A 5H7, Canada